



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

#2

REC'D 05 NOV 2003

WIPO PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02257560.9

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:  
Application no.: 02257560.9  
Demande no:

Anmeldetag:  
Date of filing: 31.10.02  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

UNILEVER PLC  
Unilever House,  
Blackfriars  
London EC4P 4BQ  
GRANDE BRETAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Water in oil emulsion comprising sterolesters

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

A23L1/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

Water in oil emulsion comprising sterolesters

Field of the invention

5 The invention relates to water in oil emulsions comprising sterol fatty acids esters and folic acid.

Background to the invention

10 Sterols are well known cholesterol lowering agents. The benefit of these ingredients to reduce the risk to cardiovascular diseases has been established for years. Where these active ingredients were initially available in the form of capsules and other pharmaceutical preparations only, over the years they  
15 have also become available in food products. The incorporation of these active ingredients in food products that are consumed daily enables the easy and reliable intake of these ingredients for many people.

20 In the search for suitable food products to incorporate sterols, it was soon found that the sterols as such are not soluble in water and show limited solubility in oil. This made the choice of food products suitable for incorporation of sterols very limited. To overcome this drawback sterols have  
25 been modified to improve their solubility in the fat phase of food products. The most common modification of sterols is to their corresponding fatty acid esters. Commercial products such as Becel pro-activ<sup>tm</sup>, and Benecol<sup>tm</sup> comprise sterol fatty acid esters.

30

EP-A-512497 and EP-A-828434 disclose food products comprising these fatty acid esters of sterols. However the products disclosed herein, especially the water in oil emulsions which

are spreadable, were found to show differences in physical properties compared to the traditional emulsions without added sterol fatty acid esters.

One of the drawbacks of addition of sterol fatty acid esters is the resulting size reducing effect on the droplet size of the dispersed aqueous phase. This size reducing effect is believed to contribute to the also signaled deterioration in mouth feel, especially in the melting properties of the water in oil emulsion. Melting of the emulsion in mouth is believed to lead to break up of the emulsion, thereby releasing the flavour and salt from the dispersed phase. A deterioration in this break up leads to a prolonged fatty impression of the emulsion and blocks the release of the flavours of the dispersed phase. Furthermore, without wishing to be bound by any theory, we

believe that a reduced dispersed phase droplet size may in certain circumstances lead to harder, sometimes more brittle products.

It is an object of the current invention to overcome at least part of these drawbacks.

#### Summary of the invention

It has now been found that the addition of folic acid to a water in oil emulsion comprising sterol fatty acid esters is effective in influencing the mouthfeel of the emulsion, the emulsion break down behaviour of the emulsion and in influencing the droplet size reducing effect of sterol fatty acid esters.

30

Therefore the present invention relates to a water in oil emulsion comprising from 30 to 90 wt% fat and from 2 to 20 wt%

of a sterol fatty acid ester, wherein the emulsion further comprises from 1.5 ppm to 1 wt% folic acid.

In a further aspect the invention relates to a method to  
5 prepare said water in oil emulsion and to use of folic acid in such emulsions.

Detailed description

Where a percentage is used, this relates to wt% unless other  
10 wise is indicated.

In the context of the invention, the terms "oil" and "fat" are used interchangeably.

15 In the context of the invention the term sterol refers to both sterols, which are also called phytosterols and to the saturated equivalents, stanols, or phytostanols.

Phytosterols, also known as plant sterols or vegetable sterols  
20 can be classified in three groups, 4-desmethylsterols, 4-monomethylsterols and 4,4'-dimethylsterols. In oils they mainly exist as free sterols and sterol esters of fatty acids although sterol glucosides and acylated sterol glucosides are also present. There are three major phytosterols namely  
25 beta-sitosterol, stigmasterol and campesterol. Schematic drawings of the components meant are as given in "Influence of Processing on Sterols of Edible Vegetable Oils", S.P. Kochhar; Prog. Lipid Res. 22: pp. 161-188.

30 The respective 5 $\alpha$ - saturated derivatives such as sitostanol, campestanol and ergostanol and their derivatives are also encompassed in the term phytosterol.

Preferably the phytosterol is selected from the group comprising fatty acid ester of  $\beta$ -sitosterol,  $\beta$ -sitostanol, campesterol, campestanol, stigmasterol, stigmastanol or a mixture thereof.

5

The phytosterols according to the invention are esterified with a fatty acid. Preferably the sterols are esterified with one or more  $C_{2-22}$  fatty acids. For the purpose of the invention the term  $C_{2-22}$  fatty acid refers to any molecule comprising a  $C_{2-22}$  main chain and at least one acid group. Although not preferred within the present context the  $C_{2-22}$  main chain may be partially substituted or side chains may be present. Preferably, however the  $C_{2-22}$  fatty acids are linear molecules comprising one or two acid group(s) as end group(s). Most preferred are linear  $C_{8-22}$  fatty acids as occur in natural oils.

Suitable examples of any such fatty acids are acetic acid, propionic acid, butyric acid, caproic acid, caprylic acid, capric acid. Other suitable acids are for example citric acid, lactic acid, oxalic acid and maleic acid. Most preferred are lauric acid, palmitic acid, stearic acid, arachidic acid, behenic acid, oleic acid, cetoleic acid, erucic acid, elaidic acid, linoleic acid and linolenic acid.

When desired a mixture of fatty acids may be used for esterification of the sterols. For example, it is possible to use a naturally occurring fat or oil as a source of the fatty acid and to carry out the esterification via an interesterification reaction.

30

In a particular embodiment, the fatty acid mixture contains a high amount (>35%, preferably >45%, further preferred >60%) of polyunsaturated fatty acids (PUFA). This does not only provide

5

the advantage of PUFA itself having good blood cholesterol lowering capacity, but also of the sterols esters prepared with such fatty acids being considered as having a higher solubility and blood cholesterol lowering efficacy in the body.

5

Preferably fatty acid mixtures of sunflower, safflower, rapeseed, linseed, linola and/or soybean are used. These are typical sources of high PUFA and/or low SAFA (saturated fatty acids). Suitable esterification conditions are for example described in WO-A-92/19640.

10

Suitable sources of phytosterols are for example soy bean, wood pulp, rapeseed oil.

15 The sterol fatty acid ester is present in an amount of from 2 to 20 wt%, preferably from 5 to 15 wt% on total emulsion weight. The preferred amount is such that the average total daily intake for a consumer taking the usual amounts of the product is from 1.6 g to 2 g free sterol per day. Hence it will  
20 be appreciated that the amount included may be dependent on the serving size of the product concerned.

The emulsions according to the invention comprise folic acid. Without wishing to be bound by any theory, it is believed that  
25 the addition of folic acid influences the mouthfeel and the way the emulsion breaks up in mouth upon consumption. This mouthfeel and emulsion break down behaviour can be judged by trained panels. It is believed that this change in mouthfeel, especially referring to melting behaviour, and emulsion  
30 breakdown characteristics is at least partly due to the influence the folic acid has on the droplet size of the dispersed aqueous phase, partly counteracting the droplet size decreasing effect of the sterolesters in the emulsion.

In the context of the invention, the term mouthfeel especially refers to the melting properties of a product on consumption. Other aspects of mouthfeel that are specifically addressed by 5 the current invention are salt and flavour release.

In the context of the invention the term folic acid includes folates and salts of folic acid such as disodium folate and folate derivatives or their salts (with for instance sodium, 10 calcium or magnesium). Folic acid, folate or derivatives thereof are preferably selected from the group comprising 5-Formyltetrahydrofolic Acid, 10-Formyltetrahydrofolic Acid, 5,10-Methenyltetrahydrofolic Acid, 5-Methyltetrahydrofolic Acid, 5,10-Methylenetetrahydrofolic Acid, Tetrahydrofolic Acid, 15 7,8-Dihydrofolic Acid, Folic Acid, Pteric Acid or combinations thereof.

Folic acid is a synthetic form of folate, a well known dietary compound. The natural sources of folate are fresh green vegetables, liver, yeast and some fruits. Our diets must daily 20 include folic acid or folate, as we cannot make it ourselves. Folic acid is part of the group of B vitamins of which also vitamin B6 and vitamin B12 are a part. While folates are both converted into an active form that the body uses, folic acid has the advantage of being more available to the body and does 25 not degrade during storage and temperature changes.

The amount of folic acid is from 1.5 ppm (0.00015 wt%) to 1 wt%, more preferred from 1.5 ppm to 0.2 wt%, even more preferred from 5 ppm (0.0005wt%) to 0.01 wt%, even more 30 preferred from 10 ppm (0.001 wt%) to 0.001 wt% on total product weight. Lower amounts were found to lead to products which did not show the desired change in mouthfeel and reduction of the size reducing effect of the sterol fatty acid esters. Higher



amounts are not recommended due to likelihood of very high intake of folic acid.

The water in oil emulsions according to the invention may be of 5 any physical texture such as pourable or spreadable. The emulsions are preferably spreadable emulsions. Even more preferred the emulsions are edible emulsions such as food products. Examples of suitable (spreadable) emulsions are margarine, butter like products, low fat spreads. Examples of 10 such compositions are disclosed in EP-A-594612, EP-A-828434.

The preferred amount of fat in these emulsions is from 30 to 60 wt%, more preferred from 35 to 45 wt%.

15 The fat may be any suitable oil or fat. Preferred fats are those known for the production of margarine and margarine derivatives such as low fat spreads. The fat is for example selected from the group comprising sunflower oil, rapeseed oil, palm oil, coconut oil, soy bean oil, palm kernel oil, butter 20 fat or a combination thereof.

Folic acid is known to participate in the metabolism of homocysteine, an amino acid in the human diet. For a number of years, high homocysteine levels have been correlated to high 25 incidence of cardiovascular disease. It is thought that lowering homocysteine may reduce the risk of cardiovascular disease. Hence the addition of folic acid to compensate for the droplet size reducing effect is beneficial for the consumer in that it simultaneously reduces homocysteine levels in blood 30 which is thought to further decrease the risk of coronary heart disease. We have surprisingly found that other B vitamins such as vitamin B6 and or vitamin B12 do not have the effect that

folic acid has on the mouthfeel, emulsion breakdown and droplet size in the presence of sterolesters in an emulsion.

Vitamins B6 and B12 are also known to interfere with the biosynthesis of purine and thiamine, to participate in the synthesis of the methyl group in the process of homocysteine methylation for producing methionine and in several growth processes. Vitamin B6 (pyridoxine hydrochloride) is a known vitamin supplement. Vitamin B12 (cyanobalamin) contributes to the health of the nervous system and is involved in the production of red blood cells. It is also known as a vitamin in food supplements.

Because of their combined positive effect on cardiovascular disease risk reduction, it is preferred that the water in oil emulsion according to the invention comprises vitamin B6 and vitamin B12 in addition to folic acid. The preferred amount of vitamin B6 is from 0.0010 to 0.5 wt% on total emulsion weight. The preferred amount of vitamin B12 is from 0.000005 to 0.0005 wt% on total emulsion weight.

The present water in oil emulsion comprises folic acid to compensate for the reduction of droplet size of the dispersed water phase. The preferred droplet size in terms of the D<sub>3,3</sub> of the dispersed water phase is from 2 to 8 µm, more preferred from 3 to 6 µm.

Optionally the emulsion comprises a thickener. Thickeners such as alginate, starch, gelatinised starch, gums including gellan gum, locust bean gum, xanthan gum or combinations thereof are commonly used to provide texture and/or viscosity in emulsion products.

Optionally the emulsion comprises an emulsifier to assist in the homogeneous admixture of aqueous phase and fat phase.

Suitable emulsifiers are preferably selected from the group comprising saturated or unsaturated monoglycerides, saturated or unsaturated diglycerides, polyoxysorbitanesters, citric acid esters, lactic acid esters or a combination thereof. Preferably emulsion comprises a monoglyceride, more preferred a saturated monoglyceride.

10 For improvement of taste and to impart nutritional value, the emulsions optionally comprise a protein. Said protein can be of vegetable or dairy origin whereby dairy origin is preferred.

Dairy proteins are suitably selected from sources such as milk protein, whey protein, skim milk powder, whey powder, sweet whey (powder), buttermilk or butter milk powder or a combination thereof. The amount of such protein is for example from 0.2 to 3, preferably from 0.5 to 2 wt% on total emulsion weight.

20 In an alternative embodiment the products comprise one or more well known antioxidants. These were found especially suitable for reduction of oxidation of fatty acids. Examples of well known antioxidants are propylgallate, tocopherol, ascorbic acid (ester).

25 Such compositions are especially suitable for use at increased pH over 5.3.

In order to further improve the oxidative stability of the products, especially of the unsaturated fatty acids contained therein, a sequestrant such as EDTA or citric acid is optionally added.

In addition to the above described ingredients the emulsions may comprise optional ingredients that are commonly used in margarine type products. Such ingredients include but are not limited to colouring agents, flavour components, preservatives  
5 such as potassium sorbate, herb, fruit or nut particles, and healthy ingredients such as vitamins, fibres.

The emulsion according to the invention may be prepared in any suitable way.

- 10 To prepare a pourable product, in an example of a suitable process a premix comprising all ingredients is prepared, followed by blending and mixing in order to establish a suitable emulsion. If desired the crystallisation of solid fat  
~~if present can either be performed a priory or as a processing~~  
15 step in which the premix is cooled by one or more scraped surface heat exchangers. In such a step also the process of emulsification could take place. Emulsification could on the other hand as well be envisaged by other kinds of techniques as, e.g. membrane emulsification and alike.

20

- For the preparation of a spreadable product, it is preferred that a premix is prepared at a temperature of about 40 to 70 °C such that the fats are melted. The composition is then preferably processed through a high shear unit with  
25 simultaneous cooling, which is optionally repeated, followed by processing in a pin stirrer and storage of the final product at a temperature from 0 to 25°C.

- We have however found that the inclusion of folic acid and  
30 optionally vitamin B6 and B12 is not straight forward because of the limited solubility of folic acid in water and the insolubility of folic acid and the vitamins in oil. This

limited solubility hinders the desired homogeneous distribution of the folic acid and the vitamins in the water in oil emulsion.

5 Therefore in a preferred process measures are taken to improve the homogeneous distribution of folic acid and the optionally added vitamins in the final water in oil emulsion.

We have surprisingly found that one or a combination of the steps (a,b,c) leads to improvement of homogeneous distribution  
10 of these ingredients throughout the emulsion.

Therefore in a preferred embodiment the invention relates to a process to prepare a water in oil emulsion comprising from 30 to 90 wt% fat and from 2 to 20 wt% of a sterol fatty acid ester, and from 1.5 ppm (0.00015 wt%) to 1 wt% folic acid  
15 wherein the folic acid is added according to any of steps (a,b,c) or a combination thereof:

- a. Folic acid is added onto a carrier and as such dosed into an emulsion or an aqueous phase
- b. Folic acid is added into the aqueous phase of an oil in  
20 water emulsion, homogenised by stirring for a few seconds, after which the emulsion is inverted into the corresponding water in oil emulsion through high speed stirring.
- c. Folic acid is pre-dispersed in one or a combination of the ingredients of the emulsion.

25

In step (a) the carrier plus folic acid is for instance made by spray drying folic acid onto maltodextrine whereby preferably the folic acid content is from 25% to 75% on total weight of the carrier and the folic acid.

30

In step (a) the emulsion or aqueous phase is preferably at 30 to 70 degrees C for better dispersion and solubility.

In step (a) the emulsion or aqueous phase are preferably stirred before and/or during the addition of the carrier with folic acid.

5

In step (b) the original oil in water emulsion comprising folic acid is the basis of the final water in oil emulsion which is obtained by inversion.

10 In step (c) the pre-dispersion can for instance be carried out in:

1. Oil with emulsifier: folic acid is added to heated oil with emulsifier whereby the oil is preferably at 50-70 °C and the emulsifier is preferably mono- and/or  
15 diglycerides at a preferred level of 0.1-1% and subsequently the oil is stirred, allowing it to homogenise for 10 minutes; or
2. Water: folic acid is added to water and the resulting dispersion is dosed into the emulsion or (part of) the  
20 aqueous phase. This pre-dispersion preferably comprises 100 to 1000 times more folic acid than the end concentration in the emulsion.
3. Sterol fatty acid esters: folic acid is added to heated  
25 sterol fatty acid esters while stirring. The temperature of the sterol ester is preferably at 60 to 70 °C. The resulting dispersion can then be added more easily to the oil phase of the emulsion.

The optional vitamin B6 and vitamin B12 are preferably added  
30 according to the same steps a, b and/or c.

13

In another aspect the invention relates to the use of folic acid to improve the mouthfeel and emulsion break down behaviour of a water in oil emulsion comprising from 30 to 90 wt% fat and from 2 to 20 wt% sterol fatty acid esters.

5

In a further aspect the invention relates to the use of folic acid to counteract the droplet size reducing effect of sterol fatty acid esters in a water in oil emulsion comprising from 30 to 90 wt% fat.

10

The invention is now illustrated in the following non-limiting examples.

## 15 Examples

### General

#### Method to determine D<sub>3,3</sub>

20 D<sub>3.3</sub> is a way to express the size (in  $\mu\text{m}$  unless otherwise stated) of a droplet of one phase/liquid in another, for instance a water droplet in oil. It is the volume weighted geometric mean water droplet diameter, i.e. 50% of the total water volume is present in droplets with a diameter larger than  
25 D<sub>3.3</sub>. To measure the D<sub>3.3</sub> Nuclear Magnetic Resonance is used. The disturbance of a magnetic field applied over an emulsion is a measure for the droplet size.

### Example 1

30

A spreadable water in oil emulsion with 35% fat (triglycerides only) was prepared with the following ingredients:

14

- Oil blend (partially hardened) rapeseed oil, sunflower oil, palm kernel oil and palm oil

35%

	• Sterol esters of beta sitosterol and sunflower oil	
5		13.9%
	• Mono/diglycerides	0.3%
	• Water	up to 100%
	• Whey powder	0.4%
	• Potassium sorbate	0.14%
10	• Alginate	0.3%
	• Maltodextrin	3.0%
	• Citric acid	0.2%
	• Beta-carotene	0.0028%

15 Folic acid was added to the water phase at 0.05 and 0.1 wt% concentrations on product. The emulsion was prepared by mixing the oil phase and water phase at 65 degrees C for a few minutes, then processing as in example 1. The tubs with the water in oil emulsion were then left to fully crystallise at 5

20 °C for 10 days.

The spreadable water in oil emulsions were evaluated by a trained sensory panel who evaluate and assess attributes of spreadable, vegetable fat based water in oil emulsions. D3,3 of the spread was measured. Main results are as follows:

25 0.05 wt% Folic acid spread: D3.3 5.37µm (sigma 2.11)  
 0.1 wt% Folic acid spread: D3.3 6.38 µm (sigma 2.11)

The salt release and mouthfeel in terms of melting behaviour of the spread containing more folic acid was perceived to be

30 better than the mouthfeel of the spread with less folic acid.



On the basis of this example it is concluded that 0.05% folic acid has a droplet size-increasing effect and that this effect is improved by increasing the folic acid level to 0.1 wt%.

5

### Example 2

Three w/o emulsions with 80% fat phase were prepared based on the following ingredients:

- 10 • Sunflower oil
  - Fully hydrogenated high erucic rapeseed oil
  - Monoglycerides
  - Sterol esters (ester of betasitosterol with sunflower oil)
  - Water
- 15 • Folic acid

#### Preparation:

- Monoglycerides, sterol esters and sunflower oil were mixed and heated up to 50°C. This fat phase makes up for 72% of the total composition, the monoglycerides account for 0.1% w/w, the
- 20 sterol esters are added at levels of 0 and 8% w/w, and sunflower oil up to 72%. After cooling down to 30°C, pre-crystallized fully hydrogenated high erucic rapeseed oil (8% w/w) in a powder form was added to the fat phase while gently stirring. After obtaining an homogeneous slurry, the water
- 25 phase was added to the fat phase while agitating with a Ultra-Turrax homogenizer at 12,900 rpm for 10 minutes. Folic acid (0.1% w/w) was pre-dispersed in the water phase prior to the addition to the fat phase. All concentrations are related to the total composition of the emulsion.
- 30 The compositions (%) and measured D3.3 and sigma of the samples (averages of three measurements) are presented in the following table 1:

Table 1; composition and results example 2

	Sample 1 (comparison)	Sample 2 (comparison)	Sample 3 (according to invention)
Sunflower oil	71.9	63.9	63.9
Fully hydrogenated high erucic rapeseed oil	8.0	8.0	8.0
Sterol esters	no	8	8
Folic acid	no	no	0.1
water	20.0	20.0	20.0
D3.3/ $\mu$ m	3.55	2.85	3.46
sigma	0.3	0.25	0.32

The data shown in the table above clearly document that the  
5 addition of sterol esters to w/o emulsions results in reduced  
average droplet sizes (approx. -20%). Further, small quantities  
of folic acid counteract this effect (approx. +21%).

## Example 3

10

Four w/o emulsions with 80% fat phase were prepared based on  
the following ingredients:

- Sunflower oil
- Fully hydrogenated high erucic rapeseed oil
- 15 • Monoglycerides
- Sterol esters (ester of betasitosterol with sunflower oil)
- Water
- Folic acid
- Vitamin B6

• Vitamin B12

Preparation:

Monoglycerides, sterolesters and sunflower oil were mixed and heated up to 50°C. This fat phase makes up for 72% of the total composition, the monoglycerides account for 0.1% w/w, the 5 sterol esters are added at levels of 8% w/w, and sunflower oil up to 72%. After cooling down to 30°C, pre-crystallized fully hydrogenated high erucic rapeseed oil (8% w/w) in a powder form was added to the fat phase while gently stirring. Folic 10 acid (0 ppm), vitamin B6 (0.05% w/w) and vitamin B12 (0.000025% w/w) were dispersed in the water phase. Then the water phase was added to the fat phase while agitating with a Ultra-Turrax homogenizer at 12,900 rpm for 10 minutes. All concentrations are related to the total composition of the emulsion.

15 The compositions (%) and measured D3.3 and sigma of the samples (averages of three measurements) are presented in the following table 2:

Table 2: composition and results example 3

	Sample 1 (comparison)	Sample 2 (according to invention)	Sample 3 (according to invention)	Sample 4 (comparison)
Sunflower oil	63.9	63.9	63.9	63.9
Fully hydrogenated high erucic rapeseed oil	8.0	8.0	8.0	8.0
Sterol esters	8	8	8	8
Folic acid	no	50 ppm	50 ppm	no
Vitamin B6	no	no	0.05	0.05
Vitamin B12	no	no	0.000025	0.000025
water	20.0	20.0	20.0	20.0
D3.3/ $\mu\text{m}$	2.85	3.06	2.99	2.89
sigma	0.25	0.31	0.38	0.22

The above results clearly show that the effect of the droplet size increase is in particular related to the presence of folic acid, and not simply due to the presence of solid material (e.g. vitamins B6 and B12) to the water phase.

*Example 4*

Two w/o emulsions with 80% fat phase were prepared with the following ingredients:

- 5 • Sunflower oil
- Fully hydrogenated high erucic rapeseed oil
- Monoglycerides
- Sterol esters (ester of betasitosterol with sunflower oil)
- Water
- 10 • Folic acid

Preparation:

Monoglycerides, sterolesters and sunflower oil were mixed and heated up to 50°C. The monoglycerides account for 0.1% w/w, the sterol esters are added at levels of 8% w/w, and sunflower oil up to 72%. After cooling down to 30°C, pre-crystallized fully hydrogenated high erucic rapeseed oil (8% w/w) in a powder form was slowly added to the fat phase while stirring. After obtaining an homogeneous slurry, the water phase was added to the fat phase while agitating with a Ultra-Turrax homogenizer at 12,900 rpm for 10 minutes. Folic acid (50 ppm) was pre-dispersed in the water phase prior to the addition to the fat phase. All concentrations are related to the total composition of the emulsion.

The compositions (%) and measured D3.3 and sigma of the samples (averages of three measurements) are presented in the following table 3:

Table 3, composition and results example 4

	Sample 1 (comparison)	Sample 2 (according to the invention)
Sunflower oil	63.9	63.9
Fully hydrogenated high erucic rapeseed oil	8.0	8.0
Sterol esters	8	8
Folic acid	no	10 ppm
water	20.0	20.0
D3.3/ $\mu$ m	2.85	3.00
sigma	0.25	0.35

It is clear from the above results that even very small quantities of folic acid have a significant effect in terms of 5 droplet size increase.

#### Example 5

Two w/o emulsions with 80% fat phase were prepared with the 10 following ingredients:

- Sunflower oil
- Fully hydrogenated high erucic rapeseed oil
- Monoglycerides
- Sterol esters (ester of betasitosterol with sunflower oil)
- 15 • Water
- Folic acid

#### Preparation:

Monoglycerides, sterolesters and sunflower oil were mixed and heated up to 50°C. The monoglycerides were added at a level of

0.1% w/w, the sterol esters at 18% w/w, and sunflower oil up to 72%. After cooling down to 30°C, pre-crystallized fully hydrogenated high erucic rapeseed oil (8% w/w) in a powder form was slowly added to the fat phase while stirring. Folic acid (0.1%) was pre-dispersed in the water phase. After obtaining an homogeneous slurry, the water phase was added to the fat phase while agitating with a Ultra-Turrax homogenizer at 12,900 rpm for 10 minutes. All concentrations are related to the total composition of the emulsion.

- 10 The compositions (%) and measured D3.3 and sigma of the samples (averages of three measurements) are presented in the following table 4:

Table 4: composition and results example 5

	Sample 1	Sample 2
Sunflower oil	53.9	53.9
Fully hydrogenated high erucic rapeseed oil	8.0	8.0
Sterol esters	18	18
Folic acid	no	0.1%
water	20.0	20.0
D3.3/ $\mu$ m	2.12	2.53
sigma	0.44	0.42

15

The results presented in the above table evidently indicate that the addition of folic acid has a significant increasing effect (approx. +19%) on the droplet size of the water phase of an emulsion with high level of sterol esters.

20

**Claims**

1. Water in oil emulsion comprising from 30 to 90 wt% fat and from 2 to 20 wt% of a sterol fatty acid ester, characterized in that the emulsion further comprises from 1.5 ppm to 1 wt % folic acid.
2. Water in oil emulsion according to claim 1 wherein the amount of folic acid is from 5 ppm to 0.01 wt%.
3. Water in oil emulsion according to claim 1 or claim 2 further comprising vitamin B6 and vitamin B12.
4. Water in oil emulsion according to claim 3 wherein the amount of vitamin B6 is from 0.0010 to 0.5 wt% on total emulsion weight and the amount of vitamin B12 is from 0.000005 to 0.0005 wt% on total emulsion weight.
5. Water in oil emulsion according to any of claims 1-4 wherein the D3,3 of the dispersed water phase is from 2 to 8  $\mu\text{m}$ .
6. Water in oil emulsion according to any of claims 1-5 wherein the sterol fatty acid ester is selected from the group comprising fatty acid ester of  $\beta$ -sitosterol,  $\beta$ -sitostanol, campesterol, campestanol, stigmasterol, stigmastanol or a mixture thereof.
7. Method for the preparation of a water in oil emulsion according to any of claims 1-6 wherein folic acid is added according to any of steps (a,b,c) or a combination thereof:
  - a. Folic acid is added onto a carrier and as such dosed into



an emulsion or an aqueous phase

b. Folic acid is added into the aqueous phase of an oil in water emulsion, homogenised by stirring for a few seconds, after which the emulsion is inverted into the corresponding water in oil emulsion through high speed stirring.

c. Folic acid is pre-dispersed in one or a combination of the ingredients of the emulsion.

8. Use of folic acid to improve the mouthfeel and emulsion break down behaviour of a water in oil emulsion comprising from 30 to 90 wt% fat and from 2 to 20 wt% sterol fatty acid esters.

**Abstract**

The invention relates to water in oil emulsions comprising a sterol fatty acid ester to which folic acid has been added to counter act the effect of the sterol fatty acid ester on the droplet size of the dispersed aqueous phase.

---

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**